



A Monthly e Magazine
ISSN:2583-2212

September 2024 Vol.4(9), 3574-3578

Popular Article

Mycotoxins in Aquafeeds

Vedant V. Kharabe¹, Ashish R. Urkude², Mitanshu Yadav³

¹M.F.Sc. Division of Aquaculture, Sher-e-Kashmir University of Agricultural Science and Technology of Kashmir (SKUAST-K), Faculty of Fisheries Rangil, Ganderbal, Jammu & Kashmir, India -190006

²M.F.Sc. Division of Aquatic Animal Health Management, Sher-e-Kashmir University of Agricultural Science and Technology of Kashmir (SKUAST-K), Faculty of Fisheries Rangil, Ganderbal, Jammu & Kashmir, India -190006

³M.F.Sc. Division of Social Sciences, Sher-e-Kashmir University of Agricultural Science and Technology of Kashmir (SKUAST-K), Faculty of Fisheries Rangil, Ganderbal, Jammu & Kashmir, India -190006

<https://doi.org/10.5281/zenodo.13768298>

Abstract

Mycotoxins are a serious problem in food and feed, particularly in fish farming where they can have detrimental effects on the fish species that are farmed. Fishmeal has been successfully substituted with plant-based ingredients in final fish diets. Nevertheless, the use of crops in feeds raises the possibility of mycotoxin and fungal contamination, as well as the frequency of mycotoxicosis in fish. Because mycotoxicosis typically causes decreased body weight, growth impairment, and greater rates of illness and mortality in fish, this could reduce aquaculture's productivity. Furthermore, the muscles of fish may accumulate certain mycotoxins. Because mycotoxins are significant genotoxins, carcinogens, and immunosuppressors to people, eating fish may therefore become another pathway for mycotoxins to infiltrate the human food chain, endangering food security and public health.

Keywords- *Mycotoxin, Immuno-suppressor, Aqua feed, Contamination, Immunity.*

1. INTRODUCTION

Fish has always been a significant dietary source of protein, and the aquaculture business is thought to be a significant contributor to the global increase in human consumption of animal food. In addition to providing protein, fish flesh has higher concentrations of important nutrients than meat from terrestrial animals, such as critical minerals, water- and fat-soluble vitamins, and trace elements. These include polyunsaturated omega-3 fatty acids (Tacon *et al.*, 2010; Tacon and Metian, 2013). At an average yearly growth rate of 10.34% (2022–2023), aquaculture has been developing and will continue to grow faster than any other significant food production sector. About 40.0% of fish raised in aquaculture for human consumption need a lot of protein rich

aquafeed that is supplied from outside sources. Both non-obligate carnivore fish, like carp, and high-trophic carnivorous fish, like salmon and tuna, which require external feed to thrive, can benefit from compound feed to promote growth and increase farmer earnings. Aquafeed is designed to not only keep fish alive but also to maximize their growth and weight gain in the least amount of time while preserving the animal's skin and muscle mass so that people can eventually devour it. It therefore depends on extremely nutrient-dense substances like fish oil and fishmeal. A significant amount of aquafeed is made up of fishmeal since it is a valuable source of micronutrients such as essential amino acids and proteins. Fish oil is a byproduct of fishmeal and a source of long-chain polyunsaturated fatty acids. Protein rich feed is essential to the productivity of aquaculture production systems, particularly intensive systems, and is thought to account for roughly 50% of overall production costs

2. MYCOTOXINS

Aspergillus, *Penicillium*, and *Fusarium* are the three genera of fungi that create mycotoxins, which are secondary metabolites that are known to be poisons and are found in almost all agricultural products and by-products globally. They have a significant negative influence on both human health and the economy. Aflatoxins (AFs), ochratoxins (OTA), fumonisins (FBs), deoxynivalenol (DON), and zearalenone (ZEA) are among the more than 400 mycotoxins that have been found (Mostrom, 2015). The danger of producing mycotoxin, particularly AFs and OTA, in aquafeed containing plant-based ingredients like wheat, corn, and soybean meal is increased during extended storage in hot and humid conditions, which promotes active fungal colonization primarily of *Aspergillus* and *Penicillium* spp.

3. MYCOTOXIN CONTAMINATION OF FISH FEED

Crops can become contaminated with mycotoxins prior to harvest, particularly if they are fortified with fibre or bran and have mold and moisture content. Inadequate storage or post-harvest conditions can also lead to contamination, as they provide an environment that is favourable for the growth of fungi and the generation of mycotoxin. Examples of such conditions include elevated temperature and increased water activity. At this time, there is no known way for getting rid of mycotoxins once they have contaminated a component or completed feed. Different processing techniques, especially those involving higher temperatures, may aid in lowering mycotoxin concentrations, though. It seems like more and more aquafeeds are including plant-based ingredients.

However, it happens almost everywhere that these components become contaminated with fungus that may be mycotoxigenic, especially *Aspergillus flavus* and *Aspergillus parasiticus*. Aflatoxin B1, ochratoxin A, and zearalenone have all been found in corn, wheat, and barley that is intended for use in fish feed in Serbia. Zearalenone levels in maize were found to be exceptionally high (mean of 5.3 mg/kg) by the authors. Low quantities of aflatoxin B1 (1.1 µg/kg to 7.4 µg/kg) were discovered in Brazil in samples of fish farm cereals, soybean bran, and maize bran.

In maize samples used for feed manufacturing, aflatoxin B1 (1.0 to 135.0 µg/kg) and fumonisins (261.0 to 2420.0 µg/kg) were also found with high incidence in Malaysia.

Aflatoxin B1 was found in all examined feed samples (soy, rice, corn, wheat, and barley) in Portugal, ranging in concentration from 1.0 to 45.0 µg/kg. Furthermore, the researchers discovered fumonisin B1 (10 to 40 µg/kg)



and deoxynivalenol (100.0 to 500.0 µg/kg) in rice, corn, wheat, and barley. All of the mycotoxins listed above that were found in feed ingredients had previously been found in aquafeed that had been completed. It is noteworthy to observe that mycotoxins are present in farm-made feed at higher concentrations and with greater frequency than in commercial feed. This may help explain why mycotoxin contamination is more widespread in developing nations because farm-made feed is more widely produced there.

All things considered, it appears that mycotoxin contamination of fish feed is a global problem, with geographic location having a significant impact on the kind and level of contamination. While, deoxynivalenol is more common in North America, Northern and Central

Europe, Africa, and North Asia, aflatoxins are most frequently found in Southern Europe, Africa, South Asia, and Southeast Asia. The most significant mycotoxins present in fish feed are discussed here, along with their effects on fish and human health. The raw ingredients of fish feed include aflatoxins, fumonisins, ochratoxins, trichothecenes, and zearalenone, in no particular order.

4. MYCOTOXINS AND THEIR EFFECTS ON FISH

1]. ALFATOXINS AND ITS PRECURSORS

After an outbreak of acute aflatoxicosis, sometimes known as "Turkey X disease," that killed about 100,000 turkeys in the 1960s led researchers to identify aflatoxins as the first mycotoxin. Aflatoxin research and characterization currently lead the group of all mycotoxins. Aflatoxin B1, B2, G1, and G2 are the most significant aflatoxins in agricultural production. *A. parasiticus* and, less frequently, *Aspergillus nomius* can also synthesize these toxins, while *A. flavus* is the primary producer of these toxins (only B-type aflatoxins). Aflatoxin-producing filamentous fungus include *Penicillium*, *Rhizopus*, *Mucor*, and *Streptomyces*.

Effects of Alfatoxins on fish:

Aflatoxin-related symptoms in fish include pale gills, poor blood coagulation, sluggish development rates, and non-increasing weight. Reduced survival rates, body darkening or yellowing, and aberrant behavior observed in young sturgeon and Nile tilapia are possible indicators of a severe infection. Fish are more vulnerable to aflatoxins depending on their age and species; that is, younger fish are more vulnerable than older fish, and certain species are more sensitive than others. The fish species most susceptible to aflatoxins is the rainbow trout (*Oncorhynchus mykiss*). Freshwater fish are said to be more susceptible to aflatoxins than warmwater fish.

2]. Fumonisin

The most dangerous fumonisin is fumonisin B1. Numerous *Fusarium* species, including *Fusarium verticillioides*, *Fusarium proliferatum*, and *Fusarium nygamai*, generate it on a regular basis. Fumonisin B1 is also produced by *Alternaria alternata*. Fumonisin is mostly found in maize and its byproducts; in Mozambique, Burkina Faso, China, and Malaysia, it has been found in 80% to 100% of corn samples.

Effects on Fish:

It appears that fumonisins cause organ damage, immune system dysfunction, decreased weight gain, and metabolic changes that raise the risk of cancer and mortality. Similar to aflatoxins, fumonisin susceptibility seems to be species-specific. It seems that channel catfish are somewhat vulnerable.



3]. Ochratoxin

Penicillium species, Aspergillus ochraceous and Aspergillus carbonarius in particular, and Aspergillus species produce ochratoxin A, the most harmful of the ochratoxins. Although it can occur in other commodities, post-harvest contamination most commonly affects cereal grains (corn, wheat, barley, and oats). Because of its extended half-life and exceptional stability, ochratoxin A is easily transferred throughout the food chain. It is also very difficult to eradicate.

Effects on Fish:

In particular, adult sea bass are extremely vulnerable to ochratoxin A. After being exposed to ochratoxin A (2.0 mg/kg and above), juvenile catfish showed a significant reduction in body weight gain within two weeks and at each subsequent weighted. Lesions in the kidney and liver were observed, and the death rate among fish fed 8.0 mg/kg of ochratoxin A rose. Feed conversion and hematocrit were also decreased.

4]. Trichothecenes

In crops like corn, wheat, barley, and oats, fungi from various genera, such as Fusarium, Myrothecium, Phomopsis, Stachybotrys, and Trichoderma, produce trichothecenes. Deoxynivalenol and T-2 toxin are the two most significant trichothecenes that are present in crops and, thus, the most harmful to animals. Despite the fact that T-2 toxin exposure affected the zebrafish embryos' health in ways such as increased mortality and malformation, cardiovascular abnormalities, and behavioural alterations.

Effects on Fish:

There is still much to learn about deoxynivalenol's effects on fish. The most delicate fish species seems to be rainbow trout in particular. This fish does not often die at a faster rate when exposed to deoxynivalenol. However, rainbow trout exposed to levels of this toxin up to 2.6 mg/kg experienced feed refusal and a decrease in feed conversion efficiency, which in turn caused a decrease in growth rate and weight gain.

5]. Zearalenone

Fusarium spp., especially *F. graminearum*, but also *Fusarium culmorum*, *Fusarium equiseti*, and *Fusarium Crookwellense*, are the main producers of zearalenone. In crops like maize, contamination by this fungus generally happens prior to harvest. According to Zhang et al. (2018), zearalenone is a mycoestrogen with significant estrogenic activity that influences the ability of many animals to reproduce.

Effects on Fish:

The ovaries of rainbow trout collected significant levels of zearalenone (up to 7.1 µg/kg), although it is yet unclear how this buildup affects reproduction. On the other hand, it has been demonstrated that brief exposure to zearalenone reduces the reproductive capacity of zebrafish (*Danio rerio*) by lowering the frequency of spawning and the relative fecundity of successive generations. Zearalenone (2.0 mg/kg), despite its immunomodulatory effects that may have compromised the fish's health, appears to boost the growth rate and feeding efficiency of rainbow trout in addition to its reproductive effects.

5. CONCLUSION

The utilization of plant-based components in aquafeed may pose significant risks to the productivity of aquaculture by increasing the rate of mycotoxin contamination. Fish that consume mycotoxins have higher sickness and death rates, more reproductive problems, less weight gain, and significant economic losses.



Additionally, even trace amounts of mycotoxins building up in fish's musculature could be extremely dangerous for health of people who eat it.

On the one hand, this adds to the already heavy burden of exposure to these harmful metabolites, especially in underdeveloped nations or areas with significant wheat consumption. Conversely, long-term exposure to low concentrations of mycotoxins in humans can lead to long-term consequences including cancer or immune system deficiencies. Therefore, the economy and public health are severely impacted by the presence of mycotoxins in aquafeed. Therefore, methods to reduce exposure and manage contamination both before and after harvest are essential. As aquaculture develops further, it should be standard procedure to monitor both the raw materials and the completed feed.

References:

- Beddington, J.R. Bioeconomic interactions between the capture fishery and aquaculture.
- Caygill, C.P.J.; Charlett, A.; Hill, M.J. Fat, fish, fish oil and cancer. *Br. J. Cancer* 1996, *Toxins* 2020.
- Deutsch, L.; Gräslund, S.; Folke, C.; Troell, M.; Huitric, M.; Kautsky, N.; Lebel, L. Feeding aquaculture growth through globalization: Exploitation of marine ecosystems for fishmeal.
- Dinter, J.; Bechthold, A.; Boeing, H.; Ellinger, S.; Leschik-Bonnet, E.; Linseisen, J.; Lorkowski, S.; Wolfram, G. Fish intake and prevention of selected nutrition-related diseases.
- FAO. Fisheries Department Aquaculture Development, Good Aquaculture Feed FAO. The State of World Fisheries and Aquaculture 2018 Meeting.
- Fernandez, E.; Chatenoud, L.; La Vecchia, C.; Negri, E.; Franceschi, S. Fish consumption and cancer risk. *Am. J. Clin.*
- Jackson, A.; Newton, R. Project to Model the Use of Fisheries by-products in the Production of Marine Ingredients with Special Reference to omega- 3 Fatty Acids EPA and DHA.
- Kobayashi, M.; Msangi, S.; Batka, M.; Vannuccini, S.; Dey, M.M.; Anderson, J.L. Fish to 2030: The Role and Opportunity for Aquaculture. *Aquac. Econ. Manag.* 2015.
- Kundam, D.N.; Acham, I.O.; Girgih, A.T. Bioactive Compounds in Fish and Their Health Benefits. *Asian Food Sci. J.* 2018.
- Manufacturing Practice; FAO: Rome, Italy, 2001.
- Mozaffarian, D.; Ascherio, A.; Hu, F.B.; Stampfer, M.J.; Willett, W.C.; Siscovick, D.S.; Rimm, E.B. Interplay Between Different Polyunsaturated Fatty Acids and Risk of Coronary Heart Disease in Men. *Circulation* 2005.
- Supartini, A.; Oishi, T.; Yagi, N. Changes in Fish Consumption Desire and Its Factors.
- Tacon, A.G.J.; Hasan, M.R.; Metian, M. Demand and Supply of Feed Ingredients for Farmed Fish and Crustaceans: Trends and Prospects; FAO: Rome, Italy, 2011.
- Tacon, A.G.J.; Metian, M. Aquaculture Feed and Food Safety: The Role of FAO and Codex Alimentarius.
- Tacon, A.G.J.; Metian, M.; Turchini, G.M.; de Silva, S.S. Responsible Aquaculture and Trophic Level Implications to Global Fish Supply.
- Tacon, A.G.J.J.; Metian, M. Fish Matters: Importance of Aquatic Foods in Human Nutrition and Global Food Supply. *Rev. Fish. Sci.* 2013.

